

MC3 BEHAVIORAL HEALTH SCREENING IN PRIMARY CARE QUICK REFERENCE

Name	Screens For	Age Group	Description	Proprietary
Developmental - American Academy of Pediatrics (AAP) recommends screening at 9, 18, 30 (or 24) months, preventative well child w/parent concern				
PEDS Parent Evaluation of Developmental Status	Parental Concerns	0-95 months	Screens for development, behavior, social emotional health – 10 items parent report	\$
ASQ3 Ages and Stages Questionnaire	Child skills and behavior	1 -60 months	Screens for developmental status of young children across five developmental areas: communication, gross motor, fine motor, problem solving, and personal-social. Parent report (6 items per each domain)	\$
Autism Spectrum Disorder – AAP recommends screening at 18, 24 months, well child visits beginning 12 mos from parent concern and observation				
M-CHAT Modified Checklist for Autism in Toddlers	Autism	16-30 months	Screening for risk of autism spectrum disorders and need for further evaluation Parent/caregiver report (20 items)	
SCQ Social Communication Questionnaire	Autism –parental report	4+years	Child developmental history and current behavior Parent/caregiver report (40 items)	\$
Psychosocial/Behavioral Assessments – AAP recommends screening at every well child visit (birth -5 years) when concerns general developmental delay				
SDQ Strengths and Difficulties Questionnaire	Psychosocial	3-17 years	Emotional, conduct, hyperactivity/inattention, peer relationship, pro-social behavior (not included in score); a separate scale assesses impact of symptoms on global functioning Parent, Teacher or self-report (25 items)	
PSC17 and PSC 35 Pediatric symptom Checklist	Psychosocial	4-16 years	General psychosocial and functional assessment in attention, externalizing and internalizing symptoms Self-administered or parent (17 or 35 items)	
ASQ-SE-2 Ages and Stages Questionnaire: Social-Emotional	Social emotional development	3-66 months	Screens children in seven areas of social-emotional development—self-regulation, compliance, social-communication, adaptive functioning, autonomy, affect, and interaction with people. Parent/caregiver report (# items vary)	\$
Depression Screening – AAP recommends 12 and older				
PHQ-A Patient Health Questionnaire-adolescent	Depression	11-18 years	Screens for depression, risk factors Self-report (9 items)	



MC3 BEHAVIORAL HEALTH SCREENING IN PRIMARY CARE QUICK REFERENCE

CDI Children's Depression Inventory	Depression	7-17 years	Screens for depression. Parent 17 items, teacher 12 items and youth 27 items, short form 10 items	\$
Substance Use (Tobacco, Alcohol, Drugs) - AAP recommends screening at 11 and up				
CRAFFT (2.1) Car, Relax, Alone, Forget, Friends, Trouble	Substance abuse	Adolescents	Screens for substance abuse (includes vaping) Self-administered or youth report (4 screener questions, then 6 items)	
Maternal Depression Screening – AAP recommends screening at 1, 2, 4, 6 months				
EDPS Edinburgh Perinatal Depression	Depression	Parent/Adult	Screens peripartum women for depression Self-report (10 items)	
PHQ-9 Patient Health Questionnaire	Depression	Adults	Screens for depression Self-report (9 items)	
Other Mental Health Screening				
ADHD				
Vanderbilt	ADHD	6-12 years	Screens in domains of inattention, disruptive behavior, anxiety, depression and separate scale functioning in school performance Parent, teacher ratings (55 items, 26 item follow up)	
Conners	ADHD	3-17 years	Assesses symptoms in opposition, cognitive, problems/inattention, hyperactivity, anxiety-shyness, perfectionism, social problems, psychosomatic problems Parent report (80 items, teacher-59 items, self-report-87 items)	\$
Anxiety				
GAD7 (General Anxiety Disorders)	Anxiety	11-17 years	Assess severity of anxiety Self-report (7 item)	
SCARED (Self-report for childhood anxiety related emotional disorders)	Anxiety	8+ years	Screens specific to anxiety-but not OCD or PTSD Parent or youth report (41 items)	
Suicide				
ASQ- Ask Suicide questions	Suicide	10-24 years	Brief screen when individuals screen positive for suicide, assesses for safety. Self-report (4 items)	



MC3 BEHAVIORAL HEALTH SCREENING IN PRIMARY CARE QUICK REFERENCE

(C-SSRS)- Columbia Suicide Severity Risk Screen	Suicide risk	5+ years	Suicide risk assessment. The answers help users identify whether someone is at risk for suicide, assess the severity and immediacy of that risk, and gauge the level of support that the person needs. Multiple forms depending on environment, interview, parent, self-report	
Trauma				
TSCC – Trauma symptom checklist for children	Trauma	3-12 years 8-16	90 item caregiver report, 54 item TSCC-A alternate does not contain sexual concern	\$
Adverse Life Experiences				
ACES –Adult Adverse Child Events Screening	Adverse life events	Adult	Stressful or traumatic events experienced before age 18. They are grouped into three categories: abuse, neglect, and household dysfunction. Self-report (10 items)	
ACE-Q-Child ACE-Q Teen	Adverse experiences	0-19	Stressful or traumatic events experienced by youth. Parent report: 17 item instrument completed by the parent/caregiver for children age 0 to 12. Teen: 19 item parent/caregiver for youth age 13 to 19 Teen/Self Report: 19 item instrument completed by youth age 13 to 19.	